VectoLex[®] WDG

Biological Larvicide

ACTIVE INGREDIENT:

Bacillus sphaericus Serotype H5a5b, Strain	
2362 dried concentrate	51.2% w/w
INERT INGREDIENTS	48.8% w/w
TOTAL	100.0% w/w

Potency: This product contains 650 BsITU/mg or 0.299 Billion BsITU/lb.

List No. 60201

EPA Reg. No. 73049-57

EPA Est. No. 33762-IA-001

INDEX:

- 1.0 Statement of Practical Treatment
- 2.0 Precautionary Statements2.1 Hazard to Humans (and Domestic Animals)2.2 Environmental Hazards
- 3.0 Directions for Use 3.1 Chemigation
- 4.0 Storage and Disposal
- 5.0 Application Directions
- 6.0 Ground and Aerial Application
- 7.0 Notice to User

KEEP OUT OF REACH OF CHILDREN CAUTION For <u>MEDICAL</u> and <u>TRANSPORT</u> Emergencies <u>ONLY</u> Call 24 Hours A Day 1-877-315-9819. For All Other Information Call 1-800-323-9597.

1.0 STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water. Call a physician if irritation persists.

2.0 PRECAUTIONARY STATEMENTS

2.1 HAZARD TO HUMANS (AND DOMESTIC ANIMALS) CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. As a general precaution when exposed to potentially high concentrations of living microbial products such as this, all mixer/loaders and applicators not in enclosed cabs or aircraft must wear a dust/mist filtering respirator meeting NIOSH standards of at least N-95, R-95, or P-95.

2.2 ENVIRONMENTAL HAZARDS

Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

3.0 DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply directly to treated finished drinking water reservoirs or drinking water receptacles.

3.1 Chemigation

Do not apply this product through any type of irrigation system.

4.0 STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in cool [59-86°F (15-30°C)], dry place.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

5.0 APPLICATION DIRECTIONS

Do not apply when area of treatment.	wind speed	d favors	drift bey	ond	the
I. For control of following non-crop		larvae	species*	in	the

Rate Range 0.5-1.5 lbs/acre (8oz-24oz/acre)		
0.5-1.5 lbs/acre (8oz-24oz/acre)		
ecies* in quito breeding occurs.		
Rate Range 0.5-1.5 lbs/acre (8oz-24oz/acre)		
Use higher rates (1 to 1.5 lbs/acre) in areas where extended residual control is necessary, or in habitats having deep water or dense surface cover.		

Apply uniformly by aerial or conventional ground equipment. Reapply as needed after 1-4 weeks.

*Mosquito species effectively controlled by VectoLex WDG:

Culex spp.	Psorophora columbiae
Aedes vexans	Psorophora ferox
Aedes melanimon	Aedes triseriatus
Aedes stimulans	Aedes sollicitans
Aedes nigromaculis	Anopheles quadrimaculatus
5	Coquillettidia perturbans

6.0 GROUND AND AERIAL APPLICATION

VectoLex WDG may be applied using conventional ground or aerial application equipment with quantities of water sufficient to provide uniform coverage of the target area. For application, first add the VectoLex WDG to water to produce a final spray mixture. The amount of water will depend on weather, spray equipment, and mosquito habitat characteristics. For application, fill the mix tank or plane hopper with the desired quantity of water. Start the mechanical or manual agitation to provide moderate circulation of water before adding the VectoLex WDG. Backpack and compressed air sprayers may be agitated by shaking after adding VectoLex WDG to the water in the sprayer. VectoLex WDG suspends readily in water and will stay suspended over normal application periods. Brief recirculation may be necessary if the spray mixture has sat for several hours or longer. Do not mix more VectoLex WDG than can be used in a 48 hour period. AVOID CONTINUOUS AGITATION OF THE SPRAY MIXTURE DURING SPRAYING. For ground spraying, apply 8-24 oz./acre (227-680 g/acre) of VectoLex WDG in 5-100 gallons of water per acre using hand-pump, airblast, mist blower, or other spray equipment. For aerial application, apply 8-24 oz./acre (227-680 g/acre) of VectoLex WDG through fixed wing or helicopter aircraft equipped with either conventional boom and nozzle systems or rotary atomizers at a convenient dilution rate. For aerial application, apply 0.5-10 gallons/acre of the final spray mixture to provide uniform coverage of the target area. Rinse and flush spray equipment thoroughly following each use.

Amount of VectoLex WDG needed to treat small areas:

	Rate of VectoLex WDG/Acre		
Area To Treat	<u>8 oz.</u>	<u>16 oz.</u>	<u>24 oz.</u>
¹ / ₄ Acre	2 oz	4 oz	6 oz
¹ / ₂ Acre	4 oz	8 oz	12 oz
10 Acres	5 lb	10 lb	15 lb
20 Acres	10 lb	20 lb	30 lb

7.0 NOTICE TO USER

SELLER MAKES NO WARRANTY, EXPRESS OR IMPLIED, OF MERCHANTABILITY, FITNESS OR OTH-ERWISE CONCERNING USE OF THIS PRODUCT OTHER THAN AS INDICATED ON THE LABEL. USER ASSUMES ALL RISKS OF USE, STORAGE OR HAN-DLING NOT IN STRICT ACCORDANCE WITH ACCOM-PANYING DIRECTIONS.

